



HAL
open science

Multi-method geochronological approach to reconstruct post-1800 floodplain sedimentation in the Upper Rhine Plain, France

Cassandra Euzen, François Chabaux, Gilles Rixhon, Frank Preusser, Frédérique Eyrolle, Valentin Chardon, Anja M Zander, Dominique Badariotti, Laurent Schmitt

► To cite this version:

Cassandra Euzen, François Chabaux, Gilles Rixhon, Frank Preusser, Frédérique Eyrolle, et al.. Multi-method geochronological approach to reconstruct post-1800 floodplain sedimentation in the Upper Rhine Plain, France. *Quaternary Geochronology*, In press, 83, pp.101561. 10.1016/j.quageo.2024.101561 . irsn-04611756

HAL Id: irsn-04611756

<https://asnr.hal.science/irsn-04611756v1>

Submitted on 25 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Innovative multi-method geochronological approach to decipher (sub-)modern floodplain sedimentation in the Upper Rhine Plain, France

5 Cassandra Euzen¹, François Chabaux², Gilles Rixhon¹, Frank Preusser³, Frédérique Eyrolle⁴, Valentin Chardon¹, Anja M. Zander⁵, Dominique Badariotti¹, Laurent Schmitt¹

- 1- Laboratoire Image Ville Environnement Université de Strasbourg CNRS ENGEEES
- 2- Institut Terre et Environnement de Strasbourg Université de Strasbourg CNRS 5 rue René Descartes
67000 Strasbourg- France
- 3- Institute of Earth and Environmental Science, Faculty of Environmental and Natural Resources,
10 University of Freiburg, Albertstr. 23b, 79104 Freiburg, Germany
- 4- Institut de Radioprotection et de Sûreté Nucléaire (IRSN) - PSE-ENV/SRTE/LRTA - Centre de Cadarache, bat 159, 13115 St Paul-lès-Durance, France
- 5- Institute of Geography, University of Cologne, Otto Fischer Str. 4, 50674 Cologne, Germany

15 *Correspondance : Cassandra Euzen (cassandra.euzen@live-cnrs.unistra.fr) François Chabaux (fchabaux@unistra.fr) Laurent Schmitt (laurent.schmitt@unistra.fr)

20 **Keywords:** Upper Rhine floodplain, geochronology, IRSL single-grain, ¹³⁷Cs, ²¹⁰Pb_{xs}, IRSL screening, portable luminescence reader, planimetric, stratigraphic and hydrological analyses, age-depth model

Abstract

The precise dating of sedimentary archives covering the last 200 years in highly human impacted alluvial floodplains is a major challenge. Combination of geochronological approaches is necessary to accurately date sub-actual sedimentary deposits. Here, we use a combination of a wide range of methods to unravel floodplain sedimentary dynamics, through the example of the Upper Rhine which is a highly engineered river. This comprises short-life radionuclides (^{137}Cs , $^{210}\text{Pb}_{\text{xs}}$), Infrared Stimulated Luminescence (IRSL) single-grain dating, as well as a global hydrogeomorphological approach (historical planimetric data, historical hydrological data, knowledge of engineering works and the morpho-sedimentary adjustments they induced). Luminescence profiling methods (IRSL screening, portable luminescence reader) were also used to further characterise sediment dynamics. Our study has demonstrated the value of old historical maps, which provide precise time markers for dating the sedimentary archive under study. We illustrate different assumptions, validity domains and limitations inherent to each method, especially the complexity of the often used ^{137}Cs to date floodplain sediments and the potential of luminescence methods for dating and estimating the sedimentation continuity. We also show the synergetic advantage of method combination in the construction of robust age models for young floodplain sedimentary archives.

1. Introduction

40 As human settlements are mainly located along rivers (Lockaby *et al.*, 2008), reconstructing the temporal evolution of their past (i) hydro-geomorphological characteristics, (ii) sediment dynamics, and (iii) water/sediment chemical composition is crucial. It not only allows understanding the human-induced impacts on their past and present functioning but also helps predicting their future evolution and building sustainable management and restoration strategies (Kondolf *et al.*, 2006; Desmet *et al.*,
45 2012). Human impacts and human-made modifications on fluvial hydrosystems increased over the last two millennia and reached huge tipping points after the onset of the Industrial Revolution in many parts of Europe, with very frequent river regulation works (Allen, 2009; Waters *et al.*, 2014; Vauclin *et al.*, 2020b). In this framework, several studies focused on monitoring suspended loads (e.g., Garzanti *et al.*, 2011; Fan *et al.*, 2020; Dendievel *et al.*, 2022) to constrain the historical releases of pollutants
50 and the human-induced impacts on hydrosystems. However, these sedimentary and/or hydrogeochemical surveys intrinsically are limited in time, starting in the second half of the 20th century at best.

Whilst the study of floodplain (including their side-channels) sediments recording periodic fluvial overbank deposition represents the best alternative for more ancient historical periods (Müller, 1977;
55 Matys Grygar *et al.*, 2012), the ability to accurately and precisely date young fluvial deposits is a crucial limitation. This is all the more true since depositional dynamics, hence sediment archival capacity, strongly varies with the spatio-temporal evolution of hydrologic connectivity between the main channel and floodplains (Dhivert *et al.*, 2015; Vauclin *et al.*, 2020a). Importantly, there is no single protocol for efficient dating of recent (<100-200 yr) fluvial deposits to date. On the one hand, classical
60 indirect approaches involve the study of historical maps (Preusser *et al.*, 2016; Hohensinner *et al.*, 2021) or historical hydrological data (Hobo *et al.*, 2010; Dhivert *et al.*, 2015; Toonen *et al.*, 2015), associated with the temporal evolution of both lateral hydrological connectivity and sediment characteristics (Eschbach *et al.*, 2018; Vauclin *et al.*, 2020b). On the other hand, both isotopic and palaeodosimetric dating methods can yield numerical ages assessing the timing of floodplain
65 deposition over the last centuries (Rixhon, 2022). In this context, the former primarily are short-life radionuclides, encompassing ¹³⁷Cs, ²⁴¹Am and ²³⁹⁺²⁴⁰Pu (Morereau, 2017; Foucher *et al.*, 2021; Klemt *et al.*, 2021), along with the method of lead excess (²¹⁰Pb_{xs}; e.g., Gobeil, 1999; Stam, 1999; Matys Grygar *et al.*, 2012; Sun *et al.*, 2019), and post-bomb radiocarbon (Ely *et al.*, 1992). Luminescence methods, originally developed to date late Pleistocene deposits (e.g., Cordier *et al.*, 2006), have been increasingly
70 applied to young fluvial environments (e.g., Madsen *et al.*, 2005; Preusser *et al.*, 2011), including the Rhine River (Preusser *et al.*, 2016; Eschbach *et al.*, 2018). Furthermore, luminescence profiling methods based on *in situ* measurements of portable readers (e.g., Sanderson and Murphy, 2010) or sample

response to test doses in the lab, so-called IRSL screening (May *et al.*, 2018). While the former were developed to identify deposition breaks or erosional phases (e.g., Muñoz-Salinas *et al.*, 2011; Kinnaird *et al.*, 2017; May *et al.*, 2018; Robins *et al.*, 2021), the latter approach provides semi-quantitative age information (e.g., Schulze *et al.*, 2022). It should be noted that the different approaches mentioned above have rarely been combined, although their combination seems very promising for improving chronological reconstructions of recent floodplain environments (Rixhon, 2022).

In this study, we foster the potential of multi-method geochronological approaches for dating recent fluvial sediments (<100-200 yr) by combining (i) relative methods based on historical maps and historical hydrological record with (ii) numerical dating using short-life radionuclide methods (^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$) and (iii) a palaeodosimetric approach, in our case single-grain infra-red stimulated luminescence (IRSL). We also use high resolution sedimentary profile scanning using a portable Optically Stimulated Luminescence (OSL) reader and lab-based IRSL screening. This large array of methods was applied to young fine-grained floodplain deposits, for which grain-size, mineralogy and chemical composition are characterized. The study site at Rhinau is located in the Upper Rhine Plain (Fig. 1 and 2), where the hydro-geomorphological adjustments induced by regulation works during the last two centuries are well known (Maire and Sanchez-Perez, 1992; Chardon *et al.*, Submitted). This innovative methodological combination aims to both (i) better grasps the limitations of each individual method and the methodological advantage of combining them and (ii) robustly constrain the history of the Rhinau sedimentary deposits.

2. Study area

The study site is located on the Rhinau Island along the Upper Rhine Plain, ~30 km upstream of the city of Strasbourg (Fig. 1). The plain represents the inner part of a major graben structure running some 350 km from SSW to NNE, from the Jura Mountains to the Rhenish Massif (e.g., Rotstein *et al.*, 2006). Up to the beginning of the 19th century, the Upper Rhine exhibited a marked downstream channel pattern evolution, with (i) a braided, (ii) a braided/anastomosing (including the Rhinau island), (iii) an anastomosing/meandering and (iv) a meandering reach (Carbiener, 1969, 1983, Fig. 1b). The functionality of the hydrosystem was heavily altered by three phases of engineering works during the 19th and 20th centuries (Honsell, 1885; Lefoulon, 1959; Schneider, 1966; Lévy, 1982; Tricart and Bravard, 1991; Descombes and Robineau, 1996; Schmitt *et al.*, 2019). The correction (AD 1840-1870) primarily aimed to control floods and to enhance navigation and agriculture. It narrowed both the braided belt within two stabilized banks spaced by 200 to 250 m and the flooding area (from 5-7 km to 1-2 km) via the construction of two high-flow dikes (Tricart and Bravard, 1991; Bernhardt, 2000; Herget *et al.*, 2005). Significant thalweg shortening resulted in a widespread average incision of 1 to

2 m over the next 150 years, peaking around 7 m at Ottmarsheim in the upstream reach (Felkel, 1969). The Rhinau area differs from this trend with a slight aggradation of ~ 0.5 m over the same timespan (Carbiener and Dillmann, 1992; Maire and Sanchez-Perez, 1992; Buck et al., 1993; Schmitt et al., 2016). The regularization (AD 1930- 1958) consisted in constructing alternate groyne fields in the main corrected channel to improve navigation (Schneider, 1966). The canalisation (AD 1928-1977) aimed to produce hydropower, develop navigation and limit floods. It mainly consisted in the construction of the so-called *Grand Canal d'Alsace* (GCA) which bypasses thanks to a diversion dam the corrected main channel, referred then to as the Old Rhine. While the GCA is continuous between Basel and Neuf-Brisach (50 km), its next downstream reach (to Strasbourg) is composed by four discontinuous sections forming artificial islands, including Rhinau. From Strasbourg to Iffezheim, the canalisation directly reworked the main corrected channel. The canalisation led to massive functional impacts, including the disconnection of (i) numerous side channels and (ii) ~ 130 km² of supplementary flooding areas (Dister et al., 1990; Schmitt et al., 2018). Finally, a large number of restoration actions such as the lowering of dikes to increase connectivity to the floodplain, have been undertaken since the 1980's (Carbiener and Schmitt, 2022; Schmitt et al., 2018).

Rhinau Island is surrounded by the GCA (constructed in AD 1963) to the West and by the by-passed Old Rhine to the East (Fig. 2a,b; Fig. 3). It is crossed by a side channel named Schaftheu, which increases the island flooding. The Old Rhine discharge is $15 \text{ m}^3 \text{ s}^{-1}$ (instream flow) when the Rhine flow measured at Basel is below $1,550 \text{ m}^3 \text{ s}^{-1}$. Over this threshold, the excess discharge is spilled into the Old Rhine by the diversion dam (Fig. 4b). Three 3.5 m-high agricultural weirs were built across the Old Rhine in AD 1963 to prevent water table (surface water and groundwater) deepening linked to the discharge drop. Consequently, the discharge threshold to flood the whole island increased from $2,000 \text{ m}^3 \text{ s}^{-1}$ to $2,650 \text{ m}^3 \text{ s}^{-1}$ after AD 1963, thereby decreasing both flood frequency and duration (Q_{md} , mean daily discharge, Maire and Sanchez-Perez, 1992; Sánchez-Pérez, 1992; Chardon et al., Submitted). Floods are induced by three processes (Maire and Sanchez-Perez, 1992): (i) bank overflows of the Old Rhine and the Schaftheu, (ii) backwater effect upstream the third agricultural weir (Fig. 2a) between AD 1963 and 2005, affecting the median section of the Schaftheu on which the study site is located (fig. 2b), and (iii) floods due to groundwater elevation above the topographical surface. Locally around the study site, the backwater effect upstream the agricultural weir compensated the Old Rhine by-passing and dewatering so the canalisation did not reduce the flood regime. Different phases of restoration works were conducted onsite (Maire et al., 1999; Meyer et al., 2013), especially in AD 2005, with the Schaftheu upstream hydrological connection enhanced (Meyer et al., 2013) (max. $Q = \sim 400 \text{ m}^3 \text{ s}^{-1}$), and interrupted the connection between the Old Rhine and the median section of the Schaftheu (except

during very high flows), dropping the backwater effect. Those restoration works probably did not
140 change significantly local flood regime since the first action roughly compensated the second.

3. Materials and methods

3.1. Field strategy: site selection and sediment sampling

Site selection to achieve the main goal of this study had two prerequisites: (i) fine-sediment deposition
starting directly after the Rhine correction and (ii) regular flooding conditions undisturbed by major
145 engineering works since then. Based on thorough analysis of both historical maps and LiDAR data, we
thus selected the site on the Rhinau island close to the Schaftheu side channel (Fig. 2a). A 1.05 m-deep
and 2 m-wide pit was dug under a forest cover (Fig. 2c) in July 2020. The stratigraphy of the fine-grained
deposits was logged onsite. Twenty-one 3 cm-thick sediment samples were collected each 5 cm for
grain-size, chemical and mineralogical analyses (Fig. 2c). Short half-life radionuclides were measured
150 in ten out of these twenty-one samples, from the subsurface (i.e., 10 cm) to a depth of 55 cm. Two
supplementary sets of twenty-one samples each were collected each 5 cm for two luminescence
profiling techniques, i.e., portable OSL reader and IRSL screening (Fig. 2c). Three samples for single-
grain IRSL dating (SG-IRSL) were collected at 76 cm, 90 cm and 105 cm depth, along with surrounding
material for dose-rate measurement. To avoid sunlight exposure, all luminescence samples were
155 collected in opaque tubes hammered in the freshly cleaned surface of the pit and sealed immediately.
In March 2021, a supplementary core was collected one meter away from the pit to enhance the
vertical resolution of sampling for short half-life radionuclides: 39 samples were collected each ~2 cm
from the surface 80 cm depth. During the 8-month apart period between pit and core sampling, a flood
occurred (~Q₂ at Basel) and generated new deposition onsite (Fig. 2,c).

3.2. Planimetric and hydrological data

160 Already successfully applied to some sections of the Upper Rhine River (e.g., Diaz-Redondo *et al.*, 2016;
Eschbach *et al.*, 2018) and its tributaries (e.g., Bruche River, Jautzy *et al.*, 2022), multi-secular
planimetric study provides insights into lateral river dynamics and geomorphological impacts induced
by engineering works. Our study is based on a set of ten successive historical co-registered maps, in
165 low flow conditions, ranging from AD 1778 to 2012, which has been manually digitalised for AD 1778,
1828, 1838, 1872, 1926 and 2012 (Fig. 3). A set of two historical aerial photographs from AD 1948 and
1976 has also been used. Hydro-morphological units separation on each digitized map is based on the
classification from Eschbach *et al.*, (2018) and Chardon *et al.* (2022b). The spatial reliability of the
digitized former maps is guaranteed by an Root Mean Square Error (RMSE) around 20 m (Chardon *et*
170 *al.*, 2022a), so it is possible to discuss changes and heterogeneities larger than 20 m. It allows for a

temporal comparison of different hydro-morphological units especially lateral evolution of the main braided belt along with the anastomosing and anabranching channels.

As no major water inflows exist between Basel, Rhinau and Rust (location in Fig. 1 and 2), flow data from Basel as well as the Rhine level in Rust are used to evaluate the evolution of these parameters on Rhinau island. Measured discharges at Basel, from AD 1830 to 2021 by the Federal Office for the Environment (FOEN, Fig. 4) as well as water-level variation between AD 1830 and 1991 at Rust (Fig. 4, from Maire and Sanchez-Perez, 1992) were combined to geomorphological data and data related to human infrastructures, to propose a reconstruction of the flooding history (Fig. 4).

3.3. Grain-size, mineralogical and geochemical analyses

After collection, sediment samples were dried at 40°C during 14 days, sieved at 2 mm and quartered with a Retsch Riffelteiler 6.5. The cumulative grain-size distribution was obtained by a Beckman Coulter laser diffraction particle size analyser. The characterization of the transport and deposition processes was based on the CM Diagram of Passega (1964, 1977), as already done on the Upper Rhine by Eschbach *et al.* (2018).

For mineralogy, bulk sediments were crushed with a Retsch agate disc mill and sieved to <125µm. Samples were mounted by lateral holder to avoid any preferred orientation of the particles. Then, analysis was conducted on a copper anticathode equipped Bruker D8 Advance instrument at 40kV and 25mA, at Institut Terre Environnement de Strasbourg (ITES) - Strasbourg - France. The angular scanning is 3-65°/2θ angular scanning every 0.014° over 0.4sec.

Analysis of major element concentrations were performed following the standard method used at ITES laboratory (e.g., Chabaux *et al.*, 2013; Rihs *et al.*, 2020) by inductively-coupled plasma atomic emission spectrometer (ICP-AES) after lithium tetraborate fusion. Samples were first crushed to <100µm and ignited in an oven successively at 110°C, 375°C (proxy of organic matter content, Heiri *et al.*, 2001) and 1000°C to evaluate sample wetness and loss on ignition (LOI). The analytical uncertainty on major element concentration was of 33 % for P₂O₅, <15 % for Na₂O, K₂O and <2 % for other major elements. Duplicates shows similar concentrations with a maximum relative difference 3 %. Measured concentration for the rock standards G-SN and BR are in good agreement with its recommended values, with less than 10 % relative difference.

3.4. ²¹⁰Pb and ¹³⁷Cs dating methods

Both sedimentary core and pit samples were analysed by gamma spectrometry. Dry samples were conditioned in 17- or 60-mL tightly closed plastic boxes for gamma counting using low-background and high-resolution Germanium Hyper pure detectors at the IRSN/LMRE laboratory in Orsay (Bouisset and

Calmet, 1997). The boxes were placed in vacuum sealed packages and stored during at least one month before measurement to ensure the secular equilibrium of the ^{210}Pb necessary to determine the concentration of $^{210}\text{Pb}_{xs}$ (^{210}Pb in excess). Efficiency calibrations were constructed using gamma-ray sources in a 1.15 g cm^{-3} density solid resin–water equivalent matrix. Activity results were corrected for true coincidence summing and self-absorption effects (Lefèvre *et al.*, 2003). Measured activities, expressed in Bq kg^{-1} dry weight are decay-corrected to the date of sampling. The activity uncertainty was estimated as the combination of calibration uncertainties, counting statistics summing and self-absorption correction uncertainties. A wide range of gamma emissions are detected with a germanium detector including ^{137}Cs , ^{210}Pb and ^{214}Bi used to determine $^{210}\text{Pb}_{xs}$. Those latter are widely used to date recent sediment deposits of the last hundred years because their half-life covers several decades: 30.08 years for the ^{137}Cs and 22.20 years for the ^{210}Pb (Goldberg, 1963).

3.4.1. ^{210}Pb dating

Lead-210 (^{210}Pb) is a daughter radionuclide of the ^{238}U radioactive decay series produced by gaseous ^{222}Rn decay. The ^{210}Pb is produced in soils and sediments where it reaches a secular equilibrium with its parents (supported ^{210}Pb) and in the atmosphere from radioactive decay of gaseous ^{222}Rn (unsupported ^{210}Pb or $^{210}\text{Pb}_{xs}$). The $^{210}\text{Pb}_{xs}$ reaches soils through dry and wet deposition from the atmosphere and is currently used for dating lake sediments where inputs of ^{210}Pb are continuous (Appleby, 2008; Foucher *et al.*, 2021). In such systems, $^{210}\text{Pb}_{xs}$ generally exponentially decreases with depth reflecting the radioactive decay of the $^{210}\text{Pb}_{xs}$ after deposition. For an evolution in closed system, the latter follows the classical radioactive decay law.

$$^{210}\text{Pb}_{xs} = ^{210}\text{Pb}_{0\ xs} e^{-\lambda t} \quad (1)$$

With $^{210}\text{Pb}_{0\ xs}$ the initial $^{210}\text{Pb}_{xs}$ at the time of deposit, λ the radioactive decay constant of ^{210}Pb (0.0311 yr^{-1}) and t the age of the deposits (e.g., in years).

By considering that the $^{210}\text{Pb}_{xs}$ flux is constant, the apparent sedimentary rate can be estimated from the linear regression of the neperian logarithm $\ln(^{210}\text{Pb}_{xs})$ versus depth (Goldberg, 1963).

$$^{210}\text{Pb}_{xs} = ^{210}\text{Pb}_{0\ xs} e^{-\lambda t} \quad (2)$$

For a constant sedimentation rate, v , equation (1) can be expressed as:

$$\ln(^{210}\text{Pb}_{xs}) = \ln(^{210}\text{Pb}_{0\ xs}) - \left(\lambda \frac{x}{v}\right) \quad (3)$$

with x the sediment depth relative to the surface (e.g. cm) and $v = \left(\frac{x}{t}\right)$ the sedimentation rate (e.g., cm yr^{-1}). In a system where the initial $^{210}\text{Pb}_{xs}$ is constant over time, equation 2 shows that depth variation of $^{210}\text{Pb}_{xs}$ in the sediment column provides the sedimentation rate v .

3.4.2. ^{137}Cs dating

235 ^{137}Cs is an anthropogenic radionuclide with a half-life of 30.2 years initially injected into the atmosphere after nuclear tests and then incorporated into soils or sediments as a result of atmospheric fallout. In western Europe two well-constrained chronological-benchmarks can be identified and used to date sedimentary records: (1) the maximum atmospheric nuclear tests in AD 1963 (Ritchie and McHenry, 1990) with worldwide fallouts and (2) the AD 1986 Chernobyl nuclear accident fallouts
240 which, unlike those of Fukushima AD 2011 (Klemt *et al.*, 2021), were generalised in Europe (Walling and He, 1997). The ^{137}Cs dating principle consists in identifying ^{137}Cs enrichment in the sediment/soil column and correlating them to specific enrichment events to infer deposition ages. As ^{137}Cs is not prone to be affected by chemical leaching or bioturbation processes (Owens *et al.*, 1996; Schimmack *et al.*, 1997; Jagercikova *et al.*, 2015), it is assumed to remain in place after deposition. The upper Rhine
245 river has been nuclearized and affected by regular releases of ^{137}Cs since the late 1960's, thereby superimposing this signal onto the atmospheric signals of AD 1963 and 1986. Here, we compare first the variations in ^{137}Cs content measured along the depth profiles (pit and core) with the known-values of the AD 1963 atmospheric fallout in the region (Renaud and Pourcelot, 2015; Meusburger *et al.*, 2020). Comparison with the AD 1986 Chernobyl fallout as well as nuclear powerplant (NPP) releases
250 that happened in the same period was achieved using (1) the ^{137}Cs values of sediment samples collected between AD 1985 and 2020 by the Institut de Radioprotection et de Sûreté Nucléaire (IRSN) along the Rhine floodplain around the Fessenheim NPP (Fig.1b) and (2) ^{137}Cs releases from Swiss NPPs (Klemt *et al.*, 2021).

3.5. Luminescence methods

255 Luminescence dating measures a latent signal that accumulates in sediment mineral grains with time, induced by the energy of ionising radiation emitted from radioactive decay (e.g., Rhodes, 2011). When exposed to light or heat stimulation, the latent signal is removed while emitting a light signal (luminescence). The strength of this signal is proportional to the amount of energy absorbed since the last resetting event and is known as the equivalent dose (D_e), which would have been required to
260 produce such a signal (in Gray = J kg^{-1}). The process of signal release is called bleaching of the signal and can be incomplete when the sediment grains are not exposed long enough to a light stimulation. After burial, ionising radiation induced by cosmogenic radiation and the radioactive decay of K, Th, and U (Rhodes, 2015) is interacting with the mineral grains at a constant energy level, known as dose rate (D , Gray per year). D_e is hence directly dependent on the dose rate (D) and the burial time.

265 Sediment dating is done measuring the D_e in quartz (Optically Stimulated Luminescence, OSL) or feldspar (Infra-Red Stimulated Luminescence, IRSL). However, Preusser *et al.* (2016) have shown for recent sediments (200 year or less) of the Upper Rhine that the OSL signal from quartz is too dim for

proper measurement and that IRSL dating must be used. These authors also demonstrated the need to use single-grains to deal with the problem of partial bleaching and this approach is hence used in the present study.

3.5.1. Single-grain IRSL Luminescence dating

The approach used here is thoroughly described in Preusser *et al.* (2016). Sample preparation was done under red-light conditions to avoid bleaching of the IRSL signal. Samples were first sieved at 150-200 μm followed by usual chemical treatments (HCl, H_2O_2 , Na-oxalate) to dissolve carbonate and break down aggregates in samples. The feldspar fraction was then isolated by density separation of the fraction $< 2.58 \text{ g cm}^{-3}$. Samples were measured using special single-grain discs with a grid of 10x10 holes, which were filled under a binocular.

For each sample, the D_e measurement, following the SAR protocol (Murray and Wintle, 2000), was performed on a Risø DA-20 (detection through 4 mm Schott BG39 and 2.1 mm Corning 7-59 glass filter), with a preheat at 150 °C for 10 s (ramp 2°s^{-1}) and stimulation for 5s using a 140mW 830 nm laser (90 % laser power) with a RG-780 filter to attenuate the second harmonic frequency (415 nm). The first 0.2s and the last 0.5s of the IR50 signal were used for D_e determination and background subtraction and the results were analysed using Analyst 4.57 (Duller, 2018) and further processed in home-made Excel sheets. For D_e determination, only grains respecting the following criteria were selected: a signal $>3\sigma$ above background, a test dose error $<15\%$, a recuperation $<10\%$, and a recycling ratio between 0.90 and 1.10 (considering uncertainties). Between 20.8 and 26.4 % of all measured grains passed the above rejection criteria and were used for D_e calculation.

For each sediment sample, a mean D_e value was calculated using both the Central Age Model (CAM) and the Minimum Age Model (MAM-3; Galbraith *et al.*, 1999). MAM-3 calculations were carried out using the R-Luminescence package (Kreutzer *et al.*, 2012; Fuchs *et al.*, 2015) applying a sigma b value of 0.20, following Preusser *et al.*, (2016). For the correction of anomalous fading (Wintle, 1973; Huntley and Lamothe, 2001), g-values were determined by storage test according to Auclair *et al.* (2003) using 2 mm multigrain discs.

The concentration of dose-rate (D) relevant elements (K, Th, U) was determined by high-resolution gamma spectrometry on the sediment surrounding 30 cm of the D_e samples, similar to the procedures described by Preusser and Kasper (2001). No evidence for radioactive disequilibrium in the Uranium decay chain was observed using the approach of Preusser and Degering (2007).

Age calculation was carried out using ADELE 2017 software (Degering and Degering, 2020), with present day geographic position and burial depth for cosmic dose rate calculation. An a-value of 0.07 ± 0.02 and an internal K-content of $12.5 \pm 1.0\%$ were used, following values reported for northern

Switzerland (Gaar *et al.*, 2014). Sediment moisture was estimated at 25 ± 5 % for the 76 cm and 90 cm samples and 30 ± 5 % for the 105 cm sample, based on sediment composition (Fig. 5) and hydrological context (§4.3).

3.5.2. Luminescence profiling: portable reader and IRSL screening

305 Samples for both methods were collected in opaque tubes in the field and under red-light conditions with removal of the outer material possibly bleached during sampling. The portable luminescence reader uses measures both IRSL and OSL net intensities from bulk samples. All samples were heated 72 hours at 60°C, manually crushed and subdivided into two aliquots to test for reproducibility. In laboratory conditions (University of Strasbourg, France), samples were successively stimulated at 880
310 ± 40 (IR) and 470 ± 20 nm (post IR-BSL) following Sanderson and Murphy's (2010) measurement protocol. The R package developed by Kreutzer *et al.* (2012) was used for post-treatment. For each aliquot the mean value of the total photon counts for the first 10 s of IRSL and post IR-OSL stimulation (minus the background value) is used.

IRSL screening measures first the natural luminescence signal emitted during IR stimulation (L_n) and
315 compares it to the sample response to a second stimulation after application of a fixed laboratory dose (T_n). Variations of the calculated L_n/T_n ratio approximates D_e as long as the signal is in the quasi-linear growing part (May *et al.*, 2018). Samples were treated first by H_2O_2 to remove organic matter content and then dried one week at 50°C. After crushing, for each sample three aliquots were prepared on steel discs and measured in a Lexsyg Smart device (Richter *et al.*, 2015) at the University of Freiburg
320 (Germany). For each aliquot the two first cycles of the SAR protocol were conducted for L_n/T_n ratio determination. The measurement includes a preheating at 250°C for 60 s, followed by IR stimulation (850 nm) at 50°C for 60 s. For the second cycle, a given test dose of ca. 22 Gy was applied with a ^{90}Sr source of 0.11 Gy s^{-1} . The ratio L_n/T_n was determined using the initial 20 s of stimulation with subtraction of the last 20 s as background of the measured IRSL signal.

325 4. Results

4.1. Geomorphological dynamics based on planimetric data

Between AD 1778 and 1828 (Fig. 3a,b), before the Rhine correction, no significant human impact affected the complex channel pattern composed by different braided belts, divided by vegetated islands, along with anabranching and anastomosing channels. The study site is located in the western
330 braided belt in AD 1778, while it is located ~300m western the braided belt in AD 1828, in the forested floodplain crossed by anastomosing and anabranching channels. As flow energy was decreasing with the distance to the braided areas, moderate-energy conditions for the study site can be proposed for

AD 1838. The Rhine correction works started locally (Fig. 3c) in AD 1838 notably, with the erection of a 1 km-long transverse dike whose western extremity is located ~400 m north-eastern the study site. Old maps indicates that during the period AD 1838-1855 (Fig.3c,d,e) the thalweg migrated westward and reached the sampling site between AD 1852 and 1855 (Fig. 3d,e). Such a migration may have resulted in the erosion of the previous fine-grained sediments (and forest). In AD 1855, the study site was located ~100 m eastern the thalweg, into the braided belt, which implies high-energy conditions with mostly coarse channel bottom sediments (gravel). During the period AD 1852-1855, the downstream thalweg curvature migration was blocked by the transverse dike, which was extended towards the west (~200 m north the study site). The artificial banks of the corrected main Rhine bed were fixed between AD 1855 and 1863 (Fig. 3e,f). The study site was in floodable conditions in AD 1863 due to the presence of local openings on the left dike of the corrected main bed, as wells as a ~1 km-long opening upstream the study site. The high-flow dike was built at around 1 km western the corrected main bed, from AD 1838 to 1863. This hydraulic configuration led to floodplain low-energy conditions, with dynamic fine sediment deposition. This also explains an important side channel narrowing, especially from AD 1855 to 1872 (Fig. 3e,f,g,h). The correction works were finished around AD 1869 (Fig. 3g), but a second high-flow dike was built from AD 1872 to 1926 (Fig. 3h,i), ~150 m western the study site. This reduced the width of the flooding area to 500 m, but probably did not modify flow energy conditions and sedimentation dynamics. The study site gradually turned, from AD 1872 to 1948 (Fig. 3h,i,j), into a forest environment associated to a decrease of the number of anastomosing channels. The AD 1948 photograph shows a moderate morphodynamic activity of the Schaftheu side channel (e.g., high variability of the channel width, flood channels close to the Schaftheu, small vegetated bars), confirming the occurrence of floods. The canalisation works induced huge planimetric changes (Fig. 3k,l), with the construction of the GCA (AD 1963) whose total width is 260 m. This explains why the upstream section of the Schaftheu was relocated eastern.

4.2. Limnimetric and hydrological evolution

The limnimetric monitoring at Rhinau shows (Fig. 4a) a mostly stable mean water level between AD 1830 and 1875, with a small lowering of about 0.25 m from 1850 to 1875. A gradual rise of about 0.5 m occurred from AD 1875 to 1930, resulting from the Rhine correction (AD 1840-1870), especially local bed aggradation due to upstream incision. The mean level remained stable until AD 1950 (nodata during AD 1938-1945). From AD 1950 to AD 1967, the mean water level decreased by 0.5 m, as a consequence of the regularization works (AD 1930-1958) and probably also of the canalisation works at Rhinau (AD 1961-1963) and Gerstheim (AD 1965-1967). Then, it abruptly raised by about 3 m in AD 1967-1968 with the backwater effect upstream the Gerstheim dam (AD 1967), which also decreased the water level fluctuation amplitude from 3 to 1-2 m.

Over the last 200 years, several intense floods ($> Q_{10}$) occurred, in AD 1852, 1876, 1881, 1882, 1910, 1999 and 2007 (Fig. 4b,c). Only three occurred since the beginning of the 20th century, but several less intense were quite relatively long (Fig. 4c). Three floods exceeding Q_{100} occurred during the 19th century, while only one occurred thereafter. The mean yearly duration of widespread flooding of the island decreased from 11 to 2 d yr⁻¹ with the canalisation (Table 1). Nevertheless, as, at the local scale of the study site the flood threshold of 2,000 m³ s⁻¹ remained roughly stable, the yearly flooding duration did not change significantly by the canalisation and certainly even increased slightly (from 11 to 14 d yr⁻¹, Table 1) due to the higher flood occurrence after the canalization than before (Fig. 4b,c).

4.3. Sedimentological, mineralogical and chemical results

In the field, macroscopic description of the sedimentary trench (e.g., colour, structure, dominant grain-size, macro-rests, roots) allowed identifying six stratigraphic units (SU; Fig. 5a) with a first gravel unit associated to sand (unit 1) below 105 cm. Unit 2 (105-92 cm), is composed by light brown-grey clayey silt (15.1 % and 70.0 % respectively). Unit 3 (92-80 cm) is brown-grey in colour and mostly composed of sand (75.7 %), with a unique mode at 213 μ m, and secondly by silt (19.8 %) (Fig. 6a). During sampling, this unit contained the groundwater table. Unit 4 (80-43 cm) is of brown-ochre-grey colour, and reflects clayey silt (18.5 % and 65.7 % respectively). It contains oxidation marks. Unit 5, is of brown-grey colour and composed of silt (71.1 %) and clay (22.4 %). The structure shows no aggregates below 33 cm, whereas some aggregates are visible above, and contains organic matter and fine roots. The dark grey-brown unit 6 is composed of clayey silt (21.3 and 72.8 % respectively) and shows many aggregates and roots.

The Passega diagram indicates that the two sandy samples of unit 3 are included in the RQ segment, indicating a graded suspension, and therefore a relatively high-energy environment (Fig. 6.b). It probably corresponds to extreme flood events after the beginning of the correction works such as the $Q_{\geq 300}$ floods of 1876 with a $Q_{md,max} = 5,530 \text{ m}^3 \text{ s}^{-1}$ and/or the floods of 1881 ($Q_{md,max} = 4,764 \text{ m}^3 \text{ s}^{-1}$, $\sim Q_{200}$) and 1882 ($Q_{md,max} = 4,371 \text{ m}^3 \text{ s}^{-1}$, $\sim Q_{70}$) (Fig. 4b). All other samples fall into the SR segment, corresponding to a uniform suspension and a low energy environment. The grain-size distributions show multiple modes, revealing a poor sediment sorting, which is common for floodplain decantation deposits. The 90 cm sample corresponds to an intermediate situation linked to a sampling bias, as it combines sediments of both units 2 and 3.

Mineralogical composition is relatively homogenous along the sedimentary column with mean composition of 32 % quartz, 31 % calcite, 18 % dolomite, 7 % albite, 3 % orthoclase, 3 % clinoclere ferroan, 3 % muscovite, 2 % montmorillonite, 1 % kaolinite and less than 1% riebeckite. The unique

exception is the sandy unit (SU3), for which the quartz content is twice as high (~57 %), while the calcite and dolomite content is twice as low (~15 % and ~10 %, respectively).

Geochemical composition and LOI_{375°C} show important variations for the sandy unit, with the highest SiO₂ (46 to 68 %) and Na₂O (0.7 to 1.2 %) concentrations, while Al₂O₃, CaO, Fe₂O₃, MgO, TiO₂, MnO, P₂O₅ and LOI_{375°C} are lower, except for K₂O. No variation from 82 cm to 52/42 cm is visible, while secondary variations exist above (Fig. 5d). This concerns Al₂O₃, K₂O, MgO, Fe₂O₃, TiO₂ and MnO, which increase between 42 and 22 cm, followed by a progressive decrease to the top. CaO, on the other hand, decreases from 16 % to 13 % between 47 and 27 cm, before increasing towards the top (15 %). Na₂O is stable from 82 cm to the top. From 52 cm to the top, SiO₂ decreases slightly and regularly (from 49 to 42 %) while P₂O₅ and LOI_{375°C} increases regularly (from 0.12 to 0.19 % and from 3 to 10 % respectively).

4.4. Short-life radionuclides

4.4.1. ¹³⁷Cs results

The figure 7a shows very similar variations with depth for both data sets. In the core the ¹³⁷Cs concentrations have <2 Bq kg⁻¹ values from the deepest sample up to ~39 cm underlying pre-bomb test ages. Above and up to 21 cm, ¹³⁷Cs increase to ~91 Bq kg⁻¹ before decreasing from 21 cm to the top where ¹³⁷Cs concentrations are at ~10 Bq kg⁻¹. This value is similar to ¹³⁷Cs concentrations measured in current Rhine sediments deposited in the Fessenheim area (~9 Bq kg⁻¹, Fig. 8). For the pit samples similar variations are observed, with a shift of ~5 cm, in comparison to the core, in the maximum ¹³⁷Cs concentration levels.

Core and pit data, taken less than 1 m away, show a global strong similarity, but differences with notably the 5 cm gap of the ¹³⁷Cs peak as well as a deposit thickness difference (101.5 cm on the core versus 105 cm on the pit) and the sandy level depth difference (73-80 cm on the core versus 80-92 cm on the pit). It is probably linked to historical higher irregular surface micro-topography than today, as well as a probable high variability of sand deposition. To compare the results of the pit (portable luminescence reader, IRSL screening, IRSL SG dating, grain-size, chemistry) and the core (²¹⁰Pb_{xs}, ¹³⁷Cs), an adjustment based on identified benchmarks (sampling dates, ¹³⁷Cs peaks and sandy level) has been done (Fig. 8a).

In the Upper Rhine area, atmospheric fallout from the AD 1986 Chernobyl NPP accident was significant and typically results in a peak concentration of ¹³⁷Cs in sedimentary records or soil profiles of these regions (Meusburger *et al.*, 2020). Monitoring of radionuclide concentrations in Rhine sediments by the French radio-protection service (IRSN) shows in particular a decrease in ¹³⁷Cs activity (decay corrected) since AD 1987. If the ¹³⁷Cs concentration peak in Rhinau pit and core is attributed to

atmospheric fallout from the Chernobyl accident our results would indicate an average sedimentation rate of $0.6 \pm 0.3 \text{ cm yr}^{-1}$ for the upper 21 cm of the core and of $0.76 \pm 0.04 \text{ cm yr}^{-1}$ for the first 26.5 centimeters of the pit (Table 2).

435 4.4.2. $^{210}\text{Pb}_{\text{xs}}$

The plot of the $^{210}\text{Pb}_{\text{xs}}$ values from the pit and the core with depth shows a good consistency between the two data sets with a linear decrease with depth (Fig.7b and Fig. 9). Interpreting that variation with the classical scheme of a closed system evolution with a same initial $^{210}\text{Pb}_{\text{xs}}$ for all the sediments at the time of their deposition (equation 2) leads to a mostly constant sedimentation rate ϵ of 0.27-0.82
440 cm yr^{-1} with a mean value of 0.41 cm yr^{-1} for the top 35 cm, on the core. With these sedimentation rates, the 35 cm depth sediments were deposited in AD 1894-1979.

4.5. Luminescence results

4.5.1. IRSL Single-Grain D and D_e values

IRSL dating on feldspar single-grain was applied on three samples collected at 76 cm, 90 cm and 105 cm
445 depth respectively. The D_e values of the individual feldspar of each sample are represented on Fig. 10a,b. The D_e scattering is, however, quite large with a more or less clear shift towards high D_e values (positive skewness). Such a shift is commonly interpreted as reflecting an incomplete resetting of some feldspar grains before their deposition, i.e. a partial bleaching of the feldspars (Murray *et al.*, 1995; Fuchs and Lang, 2001). In such a context it is recommended to calculate the age of a sample by
450 using not the D_e mean value of the feldspar population but a minimum D_e value representative of the feldspar grains without bleaching effect. These two D_e values ($D_{e,\text{CAM}}$ and $D_{e,\text{MAM}}$) are calculated by the CAM and MAM, respectively. The respective values are shown in Fig. 10a,b and the two resulting ages for each sample are given in Table 3. Determination of these ages requires the calculation of the dose rate D for each sample. The two sediment samples of 76 cm and 90 cm similar D values of 2.67 ± 0.12
455 and $2.41 \pm 0.16 \text{ Gy ka}^{-1}$, respectively, while the deeper sample has a much lower value of $1.62 \pm 0.17 \text{ Gy ka}^{-1}$ (Table 3).

4.5.2. Luminescence profiling

The results of IRSL screening and the L_x/T_x signal versus depth is plotted in Figure 7c. Variations of the L_x/T_x ratio measured for each aliquot range between 0.17 and 0.85, while mean inter-aliquot L_x/T_x ratio
460 ranges between 0.23 and 0.67. Inter-aliquots variations are small, except for 40 and 30 cm with one aliquot plotting apart from the two others. The L_x/T_x ratios in the lower half of the section (from 105 to 50 cm) increases from 0.31 to 0.62 except in the sandy layer (80-92 cm; SU3; Fig. 5a) where they are at the lowest (~ 0.23). They remain stable around 0.6 from 50 cm to 20 cm and decrease to 0.43 at the top.

465 Both IRSL and post-IR BSL profiles measured by the portable reader (in **Figure 7d**) display identical
variation trends with depth which are therefore described in the following only for the post IR-BSL
total photon counts marked by a lower standard deviation. After increasing between 95 and 90 cm to
reach 54,400 photon count at 90 cm, the post IR-BSL signal decrease between 90 and 80 cm. Then it
remains globally stable at ~14,200 photon count between 80 and 65 cm. From 65 to 45 cm, the signal
470 increases and varies significantly with two peaks of ~35,000 and 49,000 photons count at 60 and 50 cm,
respectively. These peaks, as well as the 80 cm peak, correspond to high inter-aliquots variability unlike
other samples for which aliquots have similar results. From 45 cm to the top, the post IR-BSL signal
steadily decreases from 23,400 to 10,400 photon counts. For the interpretation, it should be noted
that signal build-up depends on (i) the burial duration of the sediment, (ii) the dose rate, (iii) the
475 inherited dose at time of burial, (iv) the mineral composition and (v) the luminescence sensitivity of
the minerals (Sanderson and Murphy, 2010; Brill *et al.*, 2016).

5- Discussion

Using floodplain sedimentary deposits as environmental archive requires their accurate and reliable
dating and this can only be reached by combining complementary geochronological approaches (Hobo
480 *et al.*, 2010; Rixhon, 2022). Against this background, this study stands out because of its innovative
combination of different geomorphological and geochronological approaches. Analysis of historical
maps and aerial photographs associated with the long-term hydrological chronicles of the Rhine and
the successive engineering works, allowed us to propose a first chronologic framework for the
temporal trajectory of the studied sedimentary deposits.

485 5-1 Sedimentological evolution linked to engineering works

Historical maps (**Fig. 3**) show that the thalweg, associated with high-energy flow conditions, migrated
from east to west and flew over the study site between AD 1852 and 1855. We can assume that the
uppermost gravel unit (SU1, **Fig.5 a**) reached at a depth of 105 cm in the pit was deposited during this
short timespan (AD 1852/1855). Atop the gravel unit, the sharp grain-size boundary characterised by
490 the absence of any transitional sandy deposits (i.e. gravels overlain by silts) points to a sudden energy
drop in flow conditions and thereby well matches the first engineering works, i.e. between AD 1855-
1863. Since most of the flow was concentrated into a new bed during the Rhine correction, we can
assume fine deposition during this timespan at our sampling sites matching the 10 cm-thick silty unit
(SU2, **Fig.5a**). The 10 cm-thick sandy unit deposited atop (SU3, **Fig.5a**) bears witness of higher flow
495 conditions as attested by the Passega diagram (**Fig.6b**) which could be correlated with flooding as the
corrected riverbed would have been stabilized at this period. Two major flood events at AD 1876 and

1881-1882 (fig.4b,c) are concerned in the first place since no other major flood event happened until AD 1910.

This temporal framework, summarized on Figure 11, was completed by using independent geochronological approaches. We discuss hereafter the limits of these numerical approaches before showing the interest of combining them with the hydro-geomorphological data to constrain the history of the Rhinau sedimentary deposits.

5-2 ^{137}Cs dating in a nuclearized river

Dating sedimentary deposits by the artificial ^{137}Cs radionuclide method, emitted after AD 1950/55 is generally based on the identification of well-known ^{137}Cs concentration maxima in sediments associated to the atmospheric nuclear weapons testing maximum (1963) and the Chernobyl nuclear accident (1986) in Europe. Because of NPPs installed along the Rhine upstream of our study area, ^{137}Cs in the sediment can also be impacted by authorized ^{137}Cs liquid releases into the river. This may complicate the ^{137}Cs signal in sediments as illustrated by Eyrolle *et al.*, (2019) for the Loire River where the NPPs implanted since AD 1957 lead to increasing ^{137}Cs concentrations in the sediments before the AD 1986 Chernobyl peak. For the Aare and Rhine rivers upstream our study site, Klemm *et al.* (2021) observed five and six ^{137}Cs maxima in different sediment cores: two attributed to the classical ^{137}Cs signals from the AD 1963 and 1986 fallouts, others attributed to known ^{137}Cs releases from upstream NPPs.

By contrast, the vertical distribution of ^{137}Cs in the Rhinau's core shows only one maximum at a depth of 21 cm (Fig.8a). As top-core-sediments have similar ^{137}Cs values (10 Bq kg⁻¹, Fig.8a) to those of the Rhine sediments measured in AD 2020 by IRSN (Fig. 8b), it can be assumed that the ^{137}Cs fixed by the river sediments is directly transferred to the floodplain deposit through sedimentation processes. In the core and the pit, both the absence of bioturbation and the constant clay content - which fixes ^{137}Cs - further suggests that no secondary ^{137}Cs vertical redistribution occurred after deposition. This implicitly validates the classical interpretation that ^{137}Cs variations in the sediment column represent ^{137}Cs concentration variation at its deposition time on the sediment surface (Leenaers, 1991; Łokas *et al.*, 2010; Buchty-Lemke *et al.*, 2019; Poręba *et al.*, 2019). However, this does not explain why only one ^{137}Cs maximum is observed and does not allow age correlation (Table 2).

The onset of ^{137}Cs is usually attributed to the beginning of atmospheric nuclear weapons testing (i.e. AD 1950-55; Meybeck *et al.*, 2007; Foucher *et al.*, 2021). The ^{137}Cs concentration in sediments is relatively constant at the bottom of the core (from 80 to 40 cm) with values between 0.18 and 1.8 Bq kg⁻¹ close to the detection limits (Fig.8a). ^{137}Cs increases from 39 cm and reach a 21 cm maximum which can be correlated to the Chernobyl event based on its ^{137}Cs activity (~90 Bq kg⁻¹) which is close

530 to the ^{137}Cs activity measured for the Chernobyl peak ([70-80] Bq kg^{-1}) in the Rhine and Aare rivers by Klemt *et al.* (2021). Such an assumption implies an average sedimentation rate of 0.57-0.63 cm yr^{-1} from 0 to 21 cm and hence dates to AD 1953-1959 the beginning of ^{137}Cs concentrations increase at 39 cm depth in the core. This is consistent with the beginning of atmospheric ^{137}Cs fallouts in AD 1950-55, that implicitly supports the interpretation linking the maximum ^{137}Cs concentration at the Rhinau site to the Chernobyl peak. This age model is also reinforced by the consistency between (i) the mean sedimentation rates deduced from ^{137}Cs (0.57-0.63 cm yr^{-1} for 0-21 cm) and $^{210}\text{Pb}_{\text{xs}}$ (\in 0.27-0.82 cm yr^{-1}) and (ii) the expected ^{137}Cs value for the sediment deposited in AD 1963 and the effective ^{137}Cs value measured at Rhinau. Indeed, with a mean sedimentation rate of 0.6 cm yr^{-1} for the core, the AD 1963 level would correspond to 33.5-36 cm in the core, for which the ^{137}Cs concentration is 13 Bq kg^{-1} .
540 According to Renaud and Pourcelot (2015) and Meusburger *et al.* (2020), the atmospheric deposition of ^{137}Cs in AD 1963 reached an intensity of $\sim 3,000 \text{ Bq m}^{-2}$, and $\sim 1,200$ to $1,800 \text{ Bq kg}^{-1}$, respectively. Taking the higher value, and given our measurements were performed on a 2 to 5 cm-thick slice, whose density is 1.5-2.5 g cm^{-3} (measured onsite for some samples), deposition in AD 1963 would be [20-100] Bq kg^{-1} and thus \sim [5-25] Bq kg^{-1} measured in AD 2021.

545 This points altogether to the interpretation that the ^{137}Cs peak in the Rhinau core represents the ^{137}Cs deposition related to the Chernobyl accident in AD 1986. The absence of a clearly identified AD 1963 ^{137}Cs peak may be explained by the combination of three factors: (i) the sample thickness (2 cm on average but 2.5 and 3.5 cm for the depths corresponding to AD 1963), (ii) the relatively low sedimentation rate (0.6 cm yr^{-1}) and (iii) the occurrence of ^{137}Cs releases into the Rhine river from NPPs upstream Rhinau, i.e., Fessenheim in France (started in AD 1978) and Swiss NPPs (starting in AD 1969; Klemt *et al.*, 2021). Klemt *et al.* (2021) show significant ^{137}Cs releases into the Rhine as early as AD 1969-70. With an average sedimentation rate of 0.6 cm yr^{-1} and assimilating the ^{137}Cs peak to the Chernobyl accident, the sample corresponding to the AD 1963 level actually covers the period AD 1960-1965 and the sample just above the period AD 1965-1970. The ^{137}Cs peak therefore includes a significant contribution of ^{137}Cs released by Swiss NPPs, which explains its high ^{137}Cs activity and makes it impossible to identify the AD 1963 peak in the Rhinau sedimentary deposits. Identification of the latter would require a thinner sampling thickness than that we used - at least at this depth range. The identification of peaks for the AD 1963 nuclear fallout in the sediment profiles of the Swiss sites studied by Klemt *et al.*, (2021) was also facilitated by higher sedimentation rates (always $> 1 \text{ cm yr}^{-1}$) and higher resolution sampling.
560

The Rhinau example illustrates well that care must be taken when using ^{137}Cs for dating, and associated with $^{210}\text{Pb}_{\text{xs}}$ to infer sedimentation rates, in the case of nuclearized rivers like the Rhine with relatively low sedimentation rates. Using ^{137}Cs NPP release chronicles and/or data of ^{137}Cs monitoring in river

sediments is important to interpret the vertical ^{137}Cs variation of sedimentary archives. An alternative solution would consist in using independent tracers specific to nuclear activities, as Am-Pu, to provide complementary chronologic benchmarks (e.g., Morereau, 2017; Foucher *et al.*, 2021; Klemm *et al.*, 2021). This is, however, beyond the scope of our study.

For the top of the deposit, age-depth model (Fig.11) is precise and well-framed by ^{137}Cs which allowed calculation of two similar sedimentation rates of $\sim 0.6 \pm 0.3 \text{ cm yr}^{-1}$. Those sedimentation rates are consistent with the $^{210}\text{Pb}_{\text{xs}}$ one which is much less precise because of great uncertainties (grey dotted lines, Fig.11).

5-3 IRSL single-grain dating

The fading corrected MAM single grain ages for the two upper samples of $179 \pm 13 \text{ yr}$ (AD 1828-1854) and $163 \pm 11 \text{ yr}$ (AD 1846-1868) overlap within uncertainties and fall into the mid of the 19th century (mean estimate AD 1840-1860). This is some 30 years older (ca. 15%) than the estimate deduced from historical and hydrological data. There are different reasons that may explain a slight apparent overestimation of the SG-IRSL ages such as a presence of either small residual IRSL in the grains at the time of deposition or a minimal thermal transfer (0.04 Gy would account for the apparent offset but will be difficult to measure). Dosimetric issues related to, for example, the assumed average water content during sediment burial could also be responsible for the offset. Nevertheless, while this highlights the limitations to date very young sediment, we regard the overall age assignment as satisfying.

More pronounced is the offset between the expected basal age of the sequence, which according to historical maps should date around AD 1860. The SG-IRSL age of $243 \pm 27 \text{ yr}$ (AD 1750-1804) is significantly overestimating this estimate. However, it is striking that this much higher age is entirely explained by a much lower dose rate determined for this sample ($1.62 \pm 0.17 \text{ Gy ka}^{-1}$ for 105 cm versus $2.41 \pm 0.16 \text{ Gy ka}^{-1}$ for 92 cm), whereas the D_e is identical to the sample above. The sample RAU1_105 was taken just above the contact to underlying gravel deposits (SU1, Fig. 5a) and will almost likely have a different concentration in K, Th and U. Unfortunately, it was not considered to address this issue during fieldwork. Hence, we explain the higher basal SG-IRSL to be caused by inhomogeneity in the gamma radiation field.

5-4 Luminescence profiling techniques

Portable luminescence reader and IRSL screening methods were developed to study Quaternary sediment deposition regularity. If a sediment sequence is characterised by rather constant (i) grain size, (ii) mineralogical composition, and (iii) sedimentation rate, one can expect a gradual decrease of Ln/Tn values (May *et al.*, 2018) and portable luminescence reader signals (Portenga and Bishop, 2016)

towards the surface because of progressively shorter burial time. Any major change in sediment age would be materialized by slope breaks in Ln/Tn profiles and portable luminescence reader or (e.g. Muñoz-Salinas *et al.*, 2011; Abdulkarim *et al.*, Submitted). As no major breaks in both portable
600 luminescence reader signals are observed between 45 and 0 cm, i.e. AD 2020~1960 according to the independent ¹³⁷Cs data, this possibly points to a relatively constant sedimentation rate of silty material over this time period. This well matches our interpretation inferred from historical maps and hydrological records, suggesting a steady sedimentation at Rhinau since the Rhine canalisation (i.e. AD 1963- around the study site) and possibly slightly before.

605 However, the IRSL screening and portable luminescence reader signals do not show a joint decrease from 45 cm to the top as expected for a sedimentation of a homogeneous material at a constant rate. The data also show that all signals tend to increase from 80 to 15 cm and from 80 to 50 cm, respectively. The portable luminescence reader signals (Fig. 7d) increasing at 85 and 90 cm corresponds to major changes in the grain size with the sandy level (80-92 cm, SU3, Fig. 5a). This sandy
610 deposit probably affects the portable luminescence reader signal with (i) less bleached deposits resulting of more intense flooding events and (ii) grain size and mineralogical changes (Sanderson and Murphy, 2010; Munyikwa *et al.*, 2021). This kind of heterogeneity in recent overbank sediment deposits shows the applicability limit of the portable reader method.

The two dose rate *D* measured at 92 and 76 cm are similar (Table 3), indicating a probably low vertical
615 variation in the downer part of the profile. In the upper part, variations in the IRSL screening L_x/T_x ratio seem to correlate with variations in the concentrations of elements such as Al₂O₃, Fe₂O₃, K₂O or MnO of the sedimentary profile (Fig. 5c). Altogether this data suggests that the variation of L_x/T_x ratio the upper 50 cm of the Rhinau sediments could results from mineralogical changes and possible vertical variations of the local dose rate *D* (Sanderson and Murphy, 2010; Munyikwa *et al.*, 2021). This
620 illustrates one of the limitations for applying portable luminescence reader and IRSL screening methods for the study of depositional rates. Indeed, for short periods of deposition, portable luminescence reader or IRSL screening signals will rather reflect changes in grain size and mineralogical variations, whereas the signal increase related to the burial time might be masked.

625 5-5 Advantages of multi-method reconstruction of the age-depth model of sediment deposit

Each geochronological method used in this study reached its limit of applicability in this recent sedimentary deposition context. Nevertheless, we built an innovative robust chronometric framework by combining river planform evolution as well as stratigraphic and historical hydrological analysis with the geochronological data. The applied multi-method geochronological approach shows the advantage

630 of combining different dating tools such as old maps combined with a good knowledge of past
engineering works, grain-size and hydrology, sediment composition, and luminescence profiling
techniques (IRSL screening, portable luminescence reader), with dating methods for recent <100-200
years deposits (^{137}Cs and luminescence dating). It allows establishing robustly chronologies of the 105-
80 cm and the 40-0 cm sections of the sedimentary deposit. Old maps and hydrology based
635 chronological tags were generally confirmed by IRSL SG dating.

The obtained sedimentation rates extrapolated based on the different chronological tags (Fig.11) are
low in compared to the one obtained along the Rhone (2.9 to 7.9 cm yr⁻¹), but comparable with the
Loire ones (0.8 to 1.2 cm yr⁻¹) according to Morereau, 2017. In the upper 40 cm of the deposit, the
sedimentation rates calculated from ^{137}Cs results are of ~0.6 cm yr⁻¹ (Table 2). In detail, sedimentation
640 could be slightly accelerated on the upper 26 cm by river restauration works as shown by the
sedimentation rates obtained from the pit (Fig.11b).

Some important information is still missing on the deposition chronology between 70 and 40 cm
(Fig.11a), i.e., for the first half of the 20th century. No chronological tag could be established due to the
absence of ^{137}Cs or $^{210}\text{Pb}_{\text{xs}}$ and insufficient sands quantity for proper luminescence dating. Estimation
645 of the sedimentation rate is important because deposits resulting from floods are discontinuous.
Luminescence profiling methods could not confirm this sedimentation regularity because of the
methodological limits, certainly connected to heterogeneity of sediment deposition processes before the
canalisation. Homogeneous grain size and mineralogy suggest a relatively regular deposition and no
major changes in the sedimentation context since the beginning of the 20th century, with a mean
650 sedimentation rate of the 40-70 cm sediments similar or slightly lower than ontop of the profile
(Fig.11b). However, IRSL SG dating measured above the sandy layer suggest that part of the fine
deposits could be partly induced by the same flooding event as the sandy level, implying a much lower
sedimentation rate at the beginning of the 20th century than at the end of the 20th century.
Consequently, our results show that the canalization increased flooding around the sedimentary pit
655 and core. This change remains local as the canalisation lead to a hydraulic connectivity decrease of the
whole island of Rhinau (Maire et Sanchez-Perez, 1992; Chardon *et al.*, Submitted). This sedimentation
rate evolution at Rhinau differs from the natural filling of palaeochannels which is typically rapid at
start and decreases as the filling progresses (Malavoi and Bravard, 2011). This difference results of the
sedimentation dynamics induced by the specific Rhine engineering works.

660 Conclusions

No single method allowing to date a current sedimentary archive that goes back over the last few
centuries exist. Due to potential sedimentary discontinuities, the precise dating of floodplain archives

in strongly engineered rivers represents an important challenge along the Upper Rhine, as well as for many other rivers. The overall historical framework was defined on the basis of former maps, hydrology, isotopic (^{137}Cs , $^{210}\text{Pbxs}$) and palaeodosimetric methods (IRSL SG, portable luminescence reader, IRSL screening). We discussed advantages and limitations of both isotopic and palaeodosimetric methods in dating recent overbank sediments in a fluvial environment heavily impacted by successive engineering works over the last two centuries.

The historical maps define the beginning of the sedimentation (AD 1855-1863 here). The history of major floods generates remarkable sedimentary changes (AD 1876 inducing grain-size and mineralogical change). The IRSL SG, supports ages estimated by former maps and flood history. Although developed and optimal for dating older sediments, luminescence dating can support the deposition chronology for <200 years deposits. Finally, the upper part, post-1950, is dated with ^{137}Cs with which a sedimentation rate of $\sim 0,6 \text{ cm yr}^{-1}$ for the first 40 cm was obtained.

675 Annexe: Age-depth calculated based on ²¹⁰Pb_{xs} and ¹³⁷Cs sedimentation

rates

Figure 1: table of ages obtained by ¹³⁷Cs and ²¹⁰Pb_{xs} for each sample of the pit and the sediment core with harmonized depths. In bold are the actual samples for the core and sediment pit, with depths recalculated based on sedimentation rates obtained for the pit and core and based on the following benchmarks: end of sedimentation (2020.5 for the pit and 2021.25 for the core), base of AD 1986 peak ¹³⁷Cs and beginning of ¹³⁷Cs record (AD ~1945).

680

685

Depth		137Cs		210Pb	
pit	core	Age (mean)	Uncertainties	Age (mean)	Uncertainties
-	0,0	2021,3	-	2021,3	
-	0,5	2020,4	-	2020,0	[2019,42-2020,64]
0	0,5	2020,5	-	2020,2	[2019,60-2020,70]
-	1,5	2018,8	-	2017,6	[2015,75-2019,43]
1,5	1,6	2018,6	-	2017,3	[2015,33-2019,29]
-	2,5	2017,1	-	2015,1	[2012,08-2018,21]
-	3,5	2015,4	-	2012,7	[2008,41-2017,00]
-	4,5	2013,8	-	2010,3	[2004,74-2015,79]
6,5	5,5	2012,1	-	2007,8	[2001,11-2014,59]
-	5,5	2012,1	-	2007,8	[2001,08-2014,57]
-	6,5	2010,4	-	2005,4	[1997,41-2013,36]
-	7,5	2008,8	-	2002,9	[1993,74-2012,14]
-	8,5	2007,1	-	2000,5	[1990,07-2010,93]
11,5	9,4	2005,6	-	1998,4	[1986,89-2009,88]
-	9,5	2005,4	-	1998,1	[1986,40-2009,72]
-	11,0	2002,9	-	1994,4	[1980,90-2007,90]
-	11,5	2002,1	-	1993,2	[1979,07-2007,29]
-	13,0	1999,6	-	1989,5	[1973,56-2005,47]
16,5	13,2	1999,2	-	1988,9	[1972,67-2005,17]
-	15,0	1996,3	-	1984,6	[1966,23-2003,04]
-	16,5	1993,8	-	1981,0	[1960,73-2001,22]
-	17,0	1992,9	-	1979,8	[1958,89-2000,62]
21,5	17,1	1992,7	-	1979,5	[1958,44-2000,46]
-	19,0	1989,6	-	1974,9	[1951,56-1998,18]
26,5	21,0	1986,3	-	1970,0	[1944,22-1995,76]
-	21,5	1985,5	[1985,34-1985,6]	1968,8	[1942,39-1995,15]
-	23,0	1983,2	[1982,63-1983,65]	1965,1	[1936,88-1993,33]
-	25,0	1980,2	[1979-1981,04]	1960,2	[1929,55-1990,90]
31,5	26,2	1978,4	[1976,84-1979,49]	1957,3	[1925,17-1989,45]
-	26,5	1977,9	[1976,28-1979,09]	1956,6	[1924,05-1989,08]
-	27,0	1977,2	[1975,38-1978,44]	1955,3	[1922,21-1988,47]
-	29,0	1974,1	[1971,75-1975,83]	1950,5	[1914,88-1986,04]
36,5	31,4	1970,5	[1967,42-1972,73]	1944,6	[1906,12-1983,15]
-	31,5	1970,3	[1967,22-1972,58]	1944,4	[1905,71-1983,01]
-	31,8	1970,0	[1966,77-1972,25]	1943,7	[1904,79-1982,71]
-	34,8	1965,4	[1961,33-1968,35]	1936,4	[1893,78-1979,06]
-	36,5	1962,8	[1958,16-1966,07]	1932,2	[1887,37-1976,94]
41,5	36,6	1962,6	[1958,01-1965,96]	1932,0	[1887,07-1976,84]
-	37,0	1962,0	[1957,25-1965,42]	1930,9	[1885,53-1976,33]
-	39,0	1959,0	[1953,63-1962,81]	1926,0	[1878,19-1973,90]
-	41,0	1955,9	[1950-1960,21]	1921,2	[1870,86-1971,48]
-	41,5	1955,2	[1949,09-1959,56]	1919,9	[1869,02-1970,87]
46,5	41,8	1954,8	[1948,6-1959,2]	1919,3	[1868,02-1970,54]
51,5	47,0	1946,9	[1939,18-1952,44]	1906,6	[1848,97-1964,23]
56,5	52,2	1939,0	[1929,77-1945,68]	1893,9	[1829,92-1957,93]
61,5	57,4	1931,2	[1920,36-1938,91]	1881,2	[1810,87-1951,62]
66,5	62,5	1923,3	[1910,94-1932,15]	1868,6	[1791,82-1945,32]
71,5	67,7	1915,4	[1901,53-1925,39]	1855,9	[1772,77-1939,01]
72	68,6	1914,0	[1899,89-1924,21]	1853,7	[1769,45-1937,91]
76,5	72,9	1907,5	[1892,12-1918,63]	1843,2	[1753,72-1932,71]
81,5	78,1	1899,7	[1882,7-1911,86]	1830,5	[1734,67-1926,40]
86,5	83,3	1891,8	[1873,29-1905,1]	1817,9	[1715,62-1920,10]
91,5	88,5	1883,9	[1863,88-1898,34]	1805,2	[1696,57-1913,79]
96,5	93,7	1876,1	[1854,46-1891,58]	1792,5	[1677,52-1907,49]
101,5	98,9	1868,2	[1845,05-1884,81]	1779,8	[1658,47-1901,18]
104	101,5	1864,3	[1840,34-1881,43]	1773,5	[1648,94-1898,03]

References

- 690 ABDULKARIM, M., SCHMITT, L., FÜLLING, A., RAMBEAU, C., ERTLEN, D., MUELLER, D., ... PREUSSER, F., (Submitted), Late glacial Holocene fluvial dynamics in the Upper Rhine alluvial plain, France, *Quaternary Research*.
- ALLEN, R. C., (2009), *The British Industrial Revolution in Global Perspective*, Cambridge University Press, Cambridge.
- APPLEBY, P. G., (2008), Three decades of dating recent sediments by fallout radionuclides: a review, *The Holocene*, vol. 18, n°1, p. 83-93.
- 695 AUCLAIR, M., LAMOTHE, M. ET HUOT, S., (2003), Measurement of anomalous fading for feldspar IRSL using SAR, *Radiation Measurements*, vol. 37, n°4-5, p. 487-492.
- BERNHARDT, C., (2000), Die rheinkorrektion. Der Bürger im Staat, n°50, p. 76-81.
- BOUISSET, P. ET CALMET, D., (1997), Hyper pure gamma-ray spectrometry applied to low-level environmental sample measurements, *Geel*.
- 700 BRILL, D., JANKAEW, K. ET BRÜCKNER, H., (2016), Towards increasing the spatial resolution of luminescence chronologies – Portable luminescence reader measurements and standardized growth curves applied to a beach-ridge plain (Phra Thong, Thailand), *Quaternary Geochronology*, vol. 36, p. 134-147.
- BUCHTY-LEMKE, M., HAGEMANN, L., MAAB, A.-L., SCHÜTTRUMPF, H., SCHWARZBAUER, J. ET LEHMKUHL, F., (2019), Floodplain chronology and sedimentation rates for the past 200 years derived from trace element gradients, organic compounds, and numerical modeling, *Environmental Earth Sciences*, vol. 78, n°15, p. 445.
- 705 BUCK, W., FELKEL, K., GERHARD, H., KALWEIT, H., VAN MALDE, J., NIPPES, K.-R., ... SCHMITZ, W., (1993), *Der Rhein unter der Einwirkung des Menschen - Ausbau, Schifffahrt, Wasserwirtschaft / The Rhine under the influence of man - development, navigation, water management*, 80/261 p.
- 710 CARBIENER, R., (1969), Le Ried Centre Alsace, *Bulletin de la société industrielle de Mulhouse*, n°734.
- CARBIENER, R., (1983), Le grand Ried central d'Alsace. Ecologie et evolution d'une zone humide d'origine fluviale rhenane / The Great Central Ried of Alsace. Ecology and evolution of a wetland of Rhine fluvial origin, *Bull. Ecol.*, p. 249-277.
- 715 CARBIENER, R. ET DILLMANN, E., (1992), Cas Type de Rhinau-Daubensand: l'évolution du paysage rhéna dans la région de Rhinau, au cœur du secteur des Giessen, des Muhlbach et Brunnenwasser, In : W. A. GALLUSSER et A. SCHENKER (éd.), *Die Auen am Oberrhein / Les zones alluviales du Rhin supérieur: Ausmaß und Perspektiven des Landschaftswandels am südlichen und mittleren Oberrhein seit 1800 / Etendue et perspectives de l'évolution des paysages dans le secteur méridional et moyen du Rhin supérieur depuis 1800*, Birkhäuser, Basel, p. 113-136.
- 720 CARBIENER, R. ET SCHMITT, L., (2022), *Rhin vivant : histoire du fleuve, des poissons et des hommes*, La Nuée Bleue, Strasbourg, 287 p.
- CHABAUX, F., BLAES, E., STILLE, P., DI CHIARA ROUPERT, R., PELT, E., DOSSETO, A., ... BRANTLEY, S. L., (2013), Regolith formation rate from U-series nuclides: Implications from the study of a spheroidal weathering profile in the Rio Icacos watershed (Puerto Rico), *Geochimica et Cosmochimica Acta*, vol. 100, p. 73-95.

- 725 CHARDON, V., EUZEN, C. ET SCHMITT, L., (Submitted), Effects quantification of the Rhine regulation on the geomorphological functionality of the Rhinau-Taubergiessen alluvial forest islands and challenges for functional restoration strategies to the context of global climate change, *Journal of Environmental Management or Environmental management*.
- 730 CHARDON, V., HERRAULT, P., STAENTZEL, C., SKUPINSKI, G., FINANCE, O., WANTZEN, K. ET SCHMITT, L., (2022a), Using transition matrices to assess the spatio-temporal land cover and ecotone changes in fluvial landscapes from historical planimetric data, *Earth Surface Processes and Landforms*, p. esp.5437.
- CHARDON, V., HERRAULT, P.-A., STAENTZEL, C., FINANCE, O., SKUPINSKI, G., WANTZEN, K. M. ET SCHMITT, L., (2022b), How to assess spatio-temporal land cover and ecotones changes in fluvial landscapes from historical planimetric sources?
- 735 CORDIER, S., HARMAND, D., FRECHEN, M. ET BEINER, M., (2006), Fluvial system response to Middle and Upper Pleistocene climate change in the Meurthe and Moselle valleys (Eastern Paris Basin and Rhenish Massif), *Quaternary Science Reviews*, vol. 25, n°13, p. 1460-1474.
- DEGERING, D. ET DEGERING, A., (2020), Change is the only constant - time-dependent dose rates in luminescence dating, *Quaternary Geochronology*, vol. 58, p. 101074.
- 740 DENDIEVEL, A.-M., GROSBOIS, C., AYRAULT, S., EVRARD, O., COYNEL, A., DEBRET, M., ... MOURIER, B., (2022), Key factors influencing metal concentrations in sediments along Western European Rivers: A long-term monitoring study (1945–2020), *Science of The Total Environment*, vol. 805, p. 149778.
- DESCOMBES, R. ET ROBINEAU, G., (1996), *L'aménagement du Rhin franco-allemand / The engineering works of the Franco-German Rhine*, Société des amis du musée régional du Rhin et de la navigation.
- 745 DHIVERT, E., GROSBOIS, C., COYNEL, A., LEFÈVRE, I. ET DESMET, M., (2015), Influences of major flood sediment inputs on sedimentary and geochemical signals archived in a reservoir core (Upper Loire Basin, France), *CATENA*, vol. 126, p. 75-85.
- 750 DIAZ-REDONDO, M., EGGER, G., MARCHAMALO, M., HOHENSINNER, S. ET DISTER, E., (2016), Benchmarking Fluvial Dynamics for Process-Based River Restoration: the Upper Rhine River (1816-2014): Historical Fluvial Dynamics in the Upper Rhine, *River Research and Applications*, vol. 33, n°3, p. 403-414.
- DISTER, E., GOMER, D., OBRDLIK, P., PETERMANN, P. ET SCHNEIDER, E., (1990), Water mangement and ecological perspectives of the upper rhine's floodplains, *Regulated Rivers: Research et Management*, vol. 5, n°1, p. 1-15.
- 755 DULLER, G. A. T., (2018), Analyst v4.57 – specialist software for luminescence data, User Manual, p. 105., <https://users.aber.ac.uk/ggd/>
- ELY, L. L., WEBB, R. H. ET ENZEL, Y., (1992), Accuracy of post-bomb ¹³⁷Cs and ¹⁴C in dating fluvial deposits, *Quaternary Research*, vol. 38, n°2, p. 196-204.
- 760 ESCHBACH, D., SCHMITT, L., IMFELD, G., MAY, J.-H., PAYRAUDEAU, S., PREUSSER, F., ... SKUPINSKI, G., (2018), Long-term temporal trajectories to enhance restoration efficiency and sustainability on large rivers: an interdisciplinary study, *Hydrology and Earth System Sciences*, vol. 22, n°5, p. 2717-2737.
- EYROLLE, F., COPARD, Y., LEPAGE, H., DUCROS, L., MOREREAU, A., GROSBOIS, C., ... DESMET, M., (2019), Evidence for tritium persistence as organically bound forms in river sediments since the past nuclear weapon tests, *Scientific Reports*, vol. 9, n°1, p. 11487.

- 765 FAN, J., JIAN, X., SHANG, F., ZHANG, W., ZHANG, S. ET FU, H., (2020), Underestimated heavy metal pollution of the Minjiang River, SE China: Evidence from spatial and seasonal monitoring of suspended-load sediments, *Science of The Total Environment*, p. 142586.
- FELKEL, K., (1969), Die Erosion des Oberrheins zwischen Basel und Karlsruhe, *Gas- und Wasserfach*, n°30, p. 801-810.
- 770 FOUCHER, A., CHABOCHE, P.-A., SABATIER, P. ET EVRARD, O., (2021), A worldwide meta-analysis (1977-2020) of sediment core dating using fallout radionuclides including ^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$, *Earth System Science Data Discussions*, p. 1-61.
- FUCHS, M. C., KREUTZER, S., BUROW, C., DIETZE, M., FISCHER, M., SCHMIDT, C. ET FUCHS, M., (2015), Data processing in luminescence dating analysis: An exemplary workflow using the R package 'Luminescence', *Quaternary International*, vol. 362, p. 8-13.
- 775 FUCHS, M. ET LANG, A., (2001), OSL dating of coarse-grain fluvial quartz using single-aliquot protocols on sediments from NE Peloponnese, Greece, *Quaternary Science Reviews*, vol. 20, n°5, p. 783-787.
- GAAR, D., LOWICK, S. ET PREUSSER, F., (2014), Performance of different luminescence approaches for the dating of known-age glaciofluvial deposits from northern Switzerland, *Geochronometria*, vol. 41, n°1, p. 65-80.
- 780 GALBRAITH, R. F., ROBERTS, R. G., LASLETT, G. M., YOSHIDA, H. ET OLLEY, J. M., (1999), Optical Dating of Single and Multiple Grains of Quartz from Jinmium Rock Shelter, Northern Australia: Part I, Experimental Design and Statistical Models*, *Archaeometry*, vol. 41, n°2, p. 339-364.
- 785 GARZANTI, E., ANDÓ, S., FRANCE-LANORD, C., CENSI, P., VIGNOLA, P., GALY, V. ET LUPKER, M., (2011), Mineralogical and chemical variability of fluvial sediments 2. Suspended-load silt (Ganga–Brahmaputra, Bangladesh), *Earth and Planetary Science Letters*, vol. 302, n°1-2, p. 107-120.
- GOBEIL, C., (1999), Silver in Sediments from the St. Lawrence River and Estuary and the Saguenay Fjord, *Environmental Science et Technology*, vol. 33, n°17, p. 2953-2957.
- GOLDBERG, E. D., (1963), Geochronology with ^{210}Pb radioactive dating, International Atomic Energy Agency, Vienna, vol. 121, , p. 130.
- 790 HEIRI, O., LOTTER, A. F. ET LEMCKE, G., (2001), Loss on ignition as a method for estimating organic and carbonate content in sediments: Reproducibility and comparability of results, *Journal of Paleolimnology*, vol. 25, n°1, p. 101-110.
- HERGET, J., BREMER, E., COCH, T., DIX, A., EGGENSTEIN, G. ET EWALD, K., (2005), Engineering impact on river channels in the river Rhine catchment, p. 26.
- 795 HOBO, N., MAKASKE, B., MIDDELKOOP, H. ET WALLINGA, J., (2010), Reconstruction of floodplain sedimentation rates: a combination of methods to optimize estimates, *Earth Surface Processes and Landforms*, vol. 35, n°13, p. 1499-1515.
- HOHENSINNER, S., GRUPE, S., KLASZ, G. ET PAYER, T., (2021), Long-term deposition of fine sediments in Vienna's Danube floodplain before and after channelization, *Geomorphology*, p. 108038.
- 800 HONSELL, M., (1885), *Die Korrektion des Oberrheines : von der Schweizer Grenze unterhalb Basel bis zur Grossh. Hessischen Grenze unterhalb Mannheim : insbes. der Badische Antheil an dem Unternehmen*, Karlsruhe.

- HUNTLEY, D. J. ET LAMOTHE, M., (2001), Ubiquity of anomalous fading in K-feldspars and the measurement and correction for it in optical dating, *Canadian Journal of Earth Sciences*, vol. 38, n°7, p. 1093-1106.
- 805 JAGERCIKOVA, M., CORNU, S., LE BAS, C. ET EVRARD, O., (2015), Vertical distributions of ^{137}Cs in soils: a meta-analysis, *Journal of Soils and Sediments*, vol. 15, n°1, p. 81-95.
- JAUTZY, T., SCHMITT, L. ET RIXHON, G., (2022), Historical geomorphological adjustments of an Upper Rhine sub-tributary over the two last centuries (Bruche River, France), *Géomorphologie : Relief, Processus, Environnement*, vol. 28, n°1, p. 53-72.
- 810 KINNAIRD, T., BOLÒS, J., TURNER, A. ET TURNER, S., (2017), Optically-stimulated luminescence profiling and dating of historic agricultural terraces in Catalonia (Spain), *Journal of Archaeological Science*, vol. 78, p. 66-77.
- 815 KLEMT, E., PUTYRSKAYA, V., RÖLLIN, S., CORCHO-ALVARADO, J. A. ET SAHLI, H., (2021), Radionuclides in sediments of the Aare and Rhine river system: Fallouts, discharges, depth-age relations, mass accumulation rates and transport along the river, *Journal of Environmental Radioactivity*, vol. 232, p. 106584.
- KONDOLF, G. M., BOULTON, A. J., O'DANIEL, S., POOLE, G. C., RAHEL, F. J., STANLEY, E. H., ... NAKAMURA, K., (2006), Process-Based Ecological River Restoration: Visualizing Three-Dimensional Connectivity and Dynamic Vectors to Recover Lost Linkages, *Ecology and Society*, vol. 11, n°2, p. art5.
- 820 KREUTZER, S., SCHMIDT, C., FUCHS, M. C., DIETZE, M. ET FUCHS, M., (2012), Introducing an R package for luminescence dating analysis, vol. 30, p. 8.
- LEENAERS, H., (1991), Deposition and storage of solid-bound heavy metals in the floodplains of the River Geul (the Netherlands), *Environmental Monitoring and Assessment*, vol. 18, n°2, p. 79-103.
- 825 LEFÈVRE, O., BOUISSET, P., GERMAIN, P., BARKER, E., KERLAU, G. ET CAGNAT, X., (2003), Self-absorption correction factor applied to ^{129}I measurement by direct gamma-X spectrometry for *Fucus serratus* samples, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, vol. 506, n°1, p. 173-185.
- LEFOULON, R., (1959), L'aménagement du Rhin sur le tronçon entre Bâle et Strasbourg / Engineering works of the Rhine on the section between Basel and Strasbourg, *La Houille Blanche*, n°2, p. 203-206.
- 830 LÉVY, M., (1982), *Les aménagements du cours du Rhin au fil des siècles dans la plaine Alsace-Bade et en particulier dans la région Benfeld-Rhinou / The engineering works of the Rhine over the centuries in the Alsace-Baden plain and in particular in the Benfeld-Rhinou region.*
- LOCKABY, B. G., CONNER, W. H. ET MITCHELL, J., (2008), Floodplains, In : S. E. JØRGENSEN et B. D. FATH (éd.), *Encyclopedia of Ecology*, Academic Press, Oxford, p. 1616-1626.
- 835 ŁOKAS, E., WACHNIEW, P., CISZEWSKI, D., OWCZAREK, P. ET CHAU, N. D., (2010), Simultaneous Use of Trace Metals, ^{210}Pb and ^{137}Cs in Floodplain Sediments of a Lowland River as Indicators of Anthropogenic Impacts, *Water, Air, and Soil Pollution*, vol. 207, n°1-4, p. 57-71.
- 840 MADSEN, A. T., MURRAY, A. S., ANDERSEN, T. J., PEJRUP, M. ET BREUNING-MADSEN, H., (2005), Optically stimulated luminescence dating of young estuarine sediments: a comparison with ^{210}Pb and ^{137}Cs dating, *Marine Geology*, vol. 214, n°1-3, p. 251-268.

- MAIRE, G. ET SANCHEZ-PEREZ, J. M., (1992), Influence des aménagements hydrauliques du Rhin sur le fonctionnement hydrologique des forêts riveraines d'Alsace : l'exemple du secteur de Rhinau, *Espaces naturels rhénans - Bulletin de la société Industrielle de Mulhouse*, n°824, .
- 845 MAIRE, G., SANCHEZ-PEREZ, J.-M. ET TREMOLIERES, M., (1999), *Le réseau hydrographique de la réserve naturelle de l'île de Rhinau, fonctionnement et amélioration potentielle*, 143 p.
- MALAVOI, J.-R. ET BRAVARD, J.-P., (2011), *Éléments d'hydromorphologie fluviale. Édité par l'Onema (Office national de l'eau et des milieux aquatiques), 2010, 224 p. En ligne sur : <http://www.onema.fr/hydromorphologie-fluviale>.*
- 850 MATYS GRYGAR, T., SEDLÁČEK, J., BÁBEK, O., NOVÁKOVÁ, T., STRNAD, L. ET MIHALJEVIČ, M., (2012), Regional Contamination of Moravia (South-Eastern Czech Republic): Temporal Shift of Pb and Zn Loading in Fluvial Sediments, *Water, Air, et Soil Pollution*, vol. 223, n°2, p. 739-753.
- MAY, J.-H., MARX, S. K., REYNOLDS, W., CLARK-BALZAN, L., JACOBSEN, G. E. ET PREUSSER, F., (2018), Establishing a chronological framework for a late Quaternary seasonal swamp in the Australian 'Top End', *Quaternary Geochronology*, vol. 47, p. 81-92.
- 855 MEUSBURGER, K., EVRARD, O., ALEWELL, C., BORRELLI, P., CINELLI, G., KETTERER, M., ... BALLABIO, C., (2020), Plutonium aided reconstruction of caesium atmospheric fallout in European topsoils, *Scientific Reports*, vol. 10, n°1, p. 11858.
- 860 MEYBECK, M., LESTEL, L., BONTE, P., MOILLERON, R., COLIN, J. L., ROUSSELOT, O., ... THÉVENOT, D. R., (2007), Historical perspective of heavy metals contamination (Cd, Cr, Cu, Hg, Pb, Zn) in the Seine River basin (France) following a DPSIR approach (1950–2005), *Science of The Total Environment*, vol. 375, n°1-3, p. 204-231.
- MEYER, A., COMBROUX, I., SCHMITT, L. ET TRÉMOLIÈRES, M., (2013), Vegetation dynamics in side-channels reconnected to the Rhine River: what are the main factors controlling communities trajectories after restoration?, *Hydrobiologia*, vol. 714, n°1, p. 35-47.
- 865 MOREREAU, A., (2017), *Reconstitution à partir d'archives sédimentaires des concentrations et des sources des radionucléides ayant transité dans le Rhône et la Loire au cours de l'ère nucléaire.* (These en préparation, Aix-Marseille).
- 870 MÜLLER, G., (1977), Schadstoff-Untersuchungen an datierten Sedimentkernen aus dem Bodensee / Pollution Research on Dated Sediment Cores from Lake Constance: III. Historische Entwicklung von N- und P-Verbindungen — Beziehung zur Entwicklung von Schwermetallen und polycyclischen aromatischen Kohlenwasserstoffen /, *Zeitschrift Für Naturforschung C*, vol. 32, n°11-12, p. 920-925.
- MUÑOZ-SALINAS, E., BISHOP, P., SANDERSON, D. C. ET ZAMORANO, J.-J., (2011), Interpreting luminescence data from a portable OSL reader: three case studies in fluvial settings, *Earth Surface Processes and Landforms*, vol. 36, n°5, p. 651-660.
- 875 MUNYIKWA, K., KINNAIRD, T. C. ET SANDERSON, D. C. W., (2021), The potential of portable luminescence readers in geomorphological investigations: a review, *Earth Surface Processes and Landforms*, vol. 46, n°1, p. 131-150.
- 880 MURRAY, A. S., OLLEY, J. M. ET CAITCHEON, G. G., (1995), Measurement of equivalent doses in quartz from contemporary water-lain sediments using optically stimulated luminescence, *Quaternary Science Reviews*, vol. 14, n°4, p. 365-371.

- MURRAY, A. S. ET WINTLE, A. G., (2000), Luminescence dating of quartz using an improved single-aliquot regenerative-dose protocol, *Radiation Measurements*, vol. 32, n°1, p. 57-73.
- OWENS, P. N., WALLING, D. E. ET HE, Q., (1996), The behaviour of bomb-derived caesium-137 fallout in catchment soils, *Journal of Environmental Radioactivity*, vol. 32, n°3, p. 169-191.
- 885 PASSEGA, R., (1964), Grain Size Representation by Cm Patterns as a Geological Tool, *Journal of Sedimentary Research*, vol. 34, n°4, p. 830-847.
- PASSEGA, R., (1977), Significance of CM diagrams of sediments deposited by suspensions, *Sedimentology*, vol. 24, n°5, p. 723-733.
- 890 PORĘBA, G., ŚNIESZKO, Z., MOSKA, P., MROCZEK, P. ET MALIK, I., (2019), Interpretation of soil erosion in a Polish loess area using OSL, ^{137}Cs , $^{210}\text{Pb}_{\text{ex}}$, dendrochronology and micromorphology – case study: Biedrzykowice site (s Poland), *Geochronometria*, vol. 46, n°1, p. 57-78.
- PORTENGA, E. W. ET BISHOP, P., (2016), Confirming geomorphological interpretations based on portable OSL reader data, *Earth Surface Processes and Landforms*, vol. 41, n°3, p. 427-432.
- 895 PREUSSER, F. ET DEGERING, D., (2007), Luminescence dating of the Niederweningen mammoth site, Switzerland, *Quaternary International*, vol. 164, p. 106-112.
- PREUSSER, F. ET KASPER, H. U., (2001), Comparison of dose rate determination using high-resolution gamma spectrometry and inductively coupled plasma-mass spectrometry, *Ancient TL*, vol. 19, n°1, p. 19-23.
- 900 PREUSSER, F., MAY, J.-H., ESCHBACH, D., TRAUERSTEIN, M. ET SCHMITT, L., (2016), Infrared stimulated luminescence dating of 19th century fluvial deposits from the upper Rhine River, *Geochronometria*, vol. 43, n°1, p. 131-142.
- PREUSSER, F., SCHMITT, L., DELILE, H. ET GROSPRÊTRE, L., (2011), Optically Stimulated Luminescence (OSL) dating of the sedimentation history of the Yzeron Basin (Chaudanne sub-catchment), Rhône Valley, France, *Quaternaire*, n°vol. 22/1, p. 73-83.
- 905 RENAUD, P. ET POURCELOT, L., (2015), Les retombées en France de l'accident de Tchernobyl Rappels et situation actuelle / Fallout from the Chernobyl accident in France Background and current situation, p. 19.
- RHODES, E. J., (2011), Optically Stimulated Luminescence Dating of Sediments over the Past 200,000 Years, *Annual Review of Earth and Planetary Sciences*, vol. 39, n°1, p. 461-488.
- 910 RHODES, E. J., (2015), Dating sediments using potassium feldspar single-grain IRSL: Initial methodological considerations, *Quaternary International*, vol. 362, p. 14-22.
- RICHTER, D., RICHTER, A. ET DORNICH, K., (2015), Lexsyg smart - A luminescence detection system for dosimetry, material research and dating application, *Geochronometria*, vol. 42, n°1, p. 202-209.
- 915 RIHS, S., GONTIER, A., VOINOT, A., CHABAUX, F., PELT, E., LEMARCHAND, D. ET TURPAULT, M.-P., (2020), Field biotite weathering rate determination using U-series disequilibria, *Geochimica et Cosmochimica Acta*, vol. 276, p. 404-420.

- RITCHIE, J. C. ET MCHENRY, J. R., (1990), Application of Radioactive Fallout Cesium-137 for Measuring Soil Erosion and Sediment Accumulation Rates and Patterns: A Review, *Journal of Environmental Quality*, vol. 19, n°2, p. 215-233.
- 920 RIXHON, G., (2022), A question of time: Historical overview and modern thought on quaternary dating methods to produce fluvial chronologies, *Quaternaire. Revue de l'Association française pour l'étude du Quaternaire*, n°vol.33/2, .
- ROBINS, L., GREENBAUM, N., YU, L., BOOKMAN, R. ET ROSKIN, J., (2021), High-resolution portable-OSL analysis of Vegetated Linear Dune construction in the margins of the northwestern Negev dunefield (Israel) during the late Quaternary, *Aeolian Research*, vol. 50, p. 100680.
- 925
- ROTSTEIN, Y., EDEL, J.-B., GABRIEL, G., BOULANGER, D., SCHAMING, M. ET MUNSCHY, M., (2006), Insight into the structure of the Upper Rhine Graben and its basement from a new compilation of Bouguer Gravity, *Tectonophysics*, vol. 425, n°1-4, p. 55-70.
- SANCHEZ-PEREZ, J. M., (1992), *Fonctionnement hydrochimique d'un écosystème forestier inondable de la plaine du Rhin : la forêt alluviale du secteur de l'île de Rhinau en Alsace (France) / Hydrochemical functioning of a flooded forest ecosystem of the Rhine plain: the alluvial forest of the Rhinau island sector in Alsace (France)*.
- 930
- SANDERSON, D. C. W. ET MURPHY, S., (2010), Using simple portable OSL measurements and laboratory characterisation to help understand complex and heterogeneous sediment sequences for luminescence dating, *Quaternary Geochronology*, vol. 5, n°2-3, p. 299-305.
- 935
- SCHIMMACK, W., FLESSA, H. ET BUNZL, K., (1997), Vertical Migration of Chernobyl-Derived Radiocesium in Bavarian Grassland Soils, *Naturwissenschaften*, vol. 84, n°5, p. 204-207.
- SCHMITT, L., BEISEL, J.-N., PREUSSER, F., DE JONG, C., WANTZEN, K. M., CHARDON, V., ... BRACKHANE, S., (2019), Sustainable Management of the Upper Rhine River and Its Alluvial Plain: Lessons from Interdisciplinary Research in France and Germany, In : *Hamman Ph., Vuilleumier S., Sustainability Research in the Upper Rhine Region : Concepts and case studies, Etudes Alsaciennes et Rhénanes, Presses Universitaires de Strasbourg*, p. 201-226.
- 940
- SCHMITT, L., HOUSSIER, J., MARTIN, B., BEINER, M., SKUPINSKI, G., BOËS, E., ... OLLIVE, V., (2016), *Paléodynamique fluviale holocène dans le compartiment sud-occidental du fossé rhénan (France)*., *Revue Archéologique de l'Est, Dijon*, 403 p.
- 945
- SCHMITT, L., MORRIS, D. ET KONDOLF, G. M., (2018), Managing Floods in Large River Basins in Europe: The Rhine River, In : A. SERRA-LLOBET, G. M. KONDOLF, K. SCHAEFER et S. NICHOLSON (éd.), *Managing Flood Risk*, Springer International Publishing, Cham, p. 75-89.
- SCHNEIDER, G., (1966), *Zusammenfassende darstellung der Rheinregulierung Strassburg/Kehl-Istein. / Summary of the Rhine regulation Strasbourg/Kehl-Istein*, Freiburg, 477 p.
- 950
- SCHULZE, T., SCHWAHN, L., FÜLLING, A., ZEEDEN, C., PREUSSER, F. ET SPRAFKE, T., (2022), Investigating the loess-palaeosol sequence of Bahlingen-Schönenberg (Kaiserstuhl), southwestern Germany, using a multi-methodological approach, *Eetamp;G Quaternary Science Journal*, vol. 71, n°2, p. 145-162.
- STAM, M. H., (1999), The dating of fluvial deposits with heavy metals, ²¹⁰Pb and ¹³⁷Cs in the geul catchment (The Netherlands), *Physics and Chemistry of the Earth, Part B: Hydrology, Oceans and Atmosphere*, vol. 24, n°1-2, p. 155-160.
- 955

- SUN, X., FAN, D., LIU, M., LIAO, H. ET TIAN, Y., (2019), Persistent impact of human activities on trace metals in the Yangtze River Estuary and the East China Sea: Evidence from sedimentary records of the last 60 years, *Science of The Total Environment*, vol. 654, p. 878-889.
- 960 TOONEN, W. H. J., WINKELS, T. G., COHEN, K. M., PRINS, M. A. ET MIDDELKOOP, H., (2015), Lower Rhine historical flood magnitudes of the last 450years reproduced from grain-size measurements of flood deposits using End Member Modelling, *CATENA*, vol. 130, p. 69-81.
- TRICART, J. ET BRAVARD, J.-P., (1991), L'aménagement des trois plus grands fleuves européens : Rhin, Rhône et Danube. Problèmes et méfaits / The engineering works of the three largest European rivers: Rhine, Rhone and Danube. Problems and misdeeds, *Annales de Géographie*, vol. 100, n°561, p. 668-713.
- 965
- VAUCLIN, S., MOURIER, B., DENDIEVEL, A.-M., NOCLIN, N., PIÉGAY, H., MARCHAND, P., ... WINIARSKI, T., (2020a), Depositional environments and historical contamination as a framework to reconstruct fluvial sedimentary evolution, *Science of The Total Environment*, p. 142900.
- 970
- VAUCLIN, S., MOURIER, B., TENA, A., PIÉGAY, H. ET WINIARSKI, T., (2020b), Effects of river infrastructures on the floodplain sedimentary environment in the Rhône River, *Journal of Soils and Sediments*, vol. 20, n°6, p. 2697-2708.
- WALLING, D. E. ET HE, Q., (1997), Use of fallout ¹³⁷Cs in investigations of overbank sediment deposition on river floodplains, *CATENA*, vol. 29, n°3-4, p. 263-282.
- 975
- WATERS, C. N., ZALASIEWICZ, J. A., WILLIAMS, M., ELLIS, M. A. ET SNELLING, A. M., (2014), A stratigraphical basis for the Anthropocene?, *Geological Society, London, Special Publications*, vol. 395, n°1, p. 1-21.
- WINTLE, A. G., (1973), Anomalous Fading of Thermo-luminescence in Mineral Samples, *Nature*, vol. 245, n°5421, p. 143-144.